

Aligning Reference Frames Using Quantum States

E. Bagan, M. Baig, and R. Muñoz-Tapia

*Grup de Física Teòrica & IFAE, Facultat de Ciències, Edifici Cn,
Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona) Spain*

(Dated: February 1, 2008)

We analyze the problem of sending, in a single transmission, the information required to specify an orthogonal trihedron or reference frame through a quantum channel made out of N elementary spins. We analytically obtain the optimal strategy, i.e., the best encoding state and the best measurement. For large N , we show that the average error goes to zero linearly in $1/N$. Finally, we discuss the construction of finite optimal measurements.

PACS numbers: 03.67.Hk, 03.65.Ta

Can a system of N elementary spins be used to communicate in a single transmission the orientation of three mutually orthogonal unit vectors (orthogonal trihedron)? A positive answer would, e.g., enable two distant parties (Alice and Bob) to establish a common reference frame using just a quantum channel. This question was addressed twenty years ago by Holevo [1] who concluded that if such a quantum system has a well defined total spin J the best the sender (Alice) can attempt to achieve is to transmit the orientation of *at most one* of the three vectors. There has recently been renewed interest in this simpler, more manageable, problem of sending a single direction, and reformulations and extensions of the original question abound in the literature [2, 3, 4, 5, 6, 7, 8, 9, 10] (related issues can also be found in [11]). In all the cases, optimal communication involves collective (entangled) measurements and an accurate choice of the messenger quantum states.

In this letter, we will be concerned with the more complex problem of sending the information that specifies an orthogonal trihedron (OT). We will demonstrate that by encoding the relevant geometrical information in a particular class of states one overcomes the limitations foreseen by Holevo and a good transmission is possible. These states can be written as a simple superposition of states belonging to each of the irreducible representations of $SU(2)$ that appear in the Hilbert space of the N spins. They have maximal third component of the total spin within each representation, i.e. in standard notation are of the form $\sum C_j |j, m = j\rangle$ (therefore they are *not* eigenstates of either $\vec{J}^2$ or J_z). The quality of the optimal communication strategy is shown to increase with N and in the limit $N \rightarrow \infty$ the average error, $\langle h \rangle$, goes to zero. For large N we obtain an analytical estimate of this error, $\langle h \rangle \gtrsim 8/N$. We would like to emphasize that despite the apparent difficulty of the problem [12], an analytical treatment is possible, which provides us with a physical insight of the underlying quantum aspects involved in the communication process.

Let us suppose Alice has a system of N spins which she wishes to use to tell Bob an OT, $n = \{\vec{n}_1, \vec{n}_2, \vec{n}_3\}$. By performing quantum measurements, Bob will be able to

reconstruct this OT with some accuracy and will make the guess $n' = \{\vec{n}'_1, \vec{n}'_2, \vec{n}'_3\}$. The obvious parametrization of the different OTs is provided by the Euler angles α, β, γ , of the rotations that map $n_0 = \{\vec{x}, \vec{y}, \vec{z}\}$ into n, n' . We will use g as a shorthand for the three Euler angles, i.e., $g = (\alpha, \beta, \gamma)$. Following Holevo [1], we may quantify the quality of the communication strategy by evaluating the mean value of the error (or average error) defined for each individual measurement by

$$h(g, g') = \sum_{a=1}^3 |\vec{n}_a - \vec{n}'_a|^2 = \sum_{a=1}^3 |\vec{n}_a(g) - \vec{n}_a(g')|^2. \quad (1)$$

Assuming the OT are chosen from an isotropic distribution, and denoting by $p_{g'}(g)$ the conditional probability of Bob guessing $n(g')$ if Alice's OT is $n(g)$, one has

$$\langle h \rangle = \int dg \int dg' h(g, g') p_{g'}(g), \quad (2)$$

where dg is the Haar measure of the rotation group, $SU(2)$, which in terms of the Euler angles reads $dg = \sin \beta d\beta d\alpha d\gamma / 8\pi^2$. Covariance implies that (2) can be written as

$$\langle h \rangle = \int dg h(g, \mathbf{0}) p_{\mathbf{0}}(g), \quad (3)$$

where $\mathbf{0}$ stands for $(\alpha, \beta, \gamma) = (0, 0, 0)$. One can easily check that

$$h(g, \mathbf{0}) = 6 - 2\text{tr} U^{(1)}(g), \quad (4)$$

where $U^{(j)}$ is the $SU(2)$ irreducible representation of spin j , whose elements we write as $\mathfrak{D}_{mm'}^{(j)}(g) = \langle j, m | U^{(1)}(g) | j, m' \rangle$. One also has $t \equiv \text{tr} U^{(1)}(g) = \sum_m \mathfrak{D}_{mm}^{(1)}(g) = \cos \beta + (1 + \cos \beta) \cos(\alpha + \gamma)$. We see that the values of t lay in the real interval $[-1, 3]$. The value $t = 3$ corresponds to perfect determination of Alice's OT and implies that $\langle h \rangle = 0$. Note also that $\langle h \rangle = 6 - 2\langle t \rangle$. Random guessing implies $\langle t \rangle = 0$ ($\langle h \rangle = 6$), while perfect determination of one axis and random guessing of the remaining two imply $\langle t \rangle = 1$ ($\langle h \rangle = 4$).

The most general quantum state Alice can use has the form $|A(g)\rangle = U(g)|A\rangle$. Here $U(g) = \bigoplus_j U^{(j)}$ and

$$|A\rangle = \sum_j |A^j\rangle = \sum_{j,m} A_m^j |j, m\rangle; \quad \sum_{j,m} |A_m^j|^2 = 1, \quad (5)$$

where j runs from 0 to $N/2$ (for simplicity we will only consider N even) and m runs from $-j$ to j . $|A\rangle$ is a fixed reference state associated with the OT n_0 .

Likewise, we may write a reference state $|B\rangle$ from which we can construct Bob's projectors of his Positive Operator Valued Measurement (POVM). The general form of the state is

$$|B\rangle = \sum_j \sqrt{2j+1} |B^j\rangle; \quad |B^j\rangle = \sum_m B_m^j |j, m\rangle, \quad (6)$$

where the square root is introduced for later convenience, and the projectors are

$$O(g) = U(g)|B\rangle\langle B|U^\dagger(g). \quad (7)$$

We will first consider continuum POVMs for simplicity but finite ones can also be constructed, as will be explained below. The condition $\mathbb{I} = \int dg O(g)$ requires that

$$\sum_{m=-j}^j |B_m^j|^2 = 1, \quad \forall j, \quad (8)$$

as can be easily shown with the help of the orthogonality relations

$$\int dg \mathfrak{D}_{Mm}^{(J)}(g) \mathfrak{D}_{M'm'}^{(J')*}(g) = \frac{\delta^{JJ'} \delta_{MM'} \delta_{mm'}}{2J+1}. \quad (9)$$

Quantum Mechanics tells us that $p_0(g) = |\langle B|U(g)|A\rangle|^2$, hence we have

$$\langle t \rangle = \int dg |\langle B|U(g)|A\rangle|^2 \text{tr} U^{(1)}. \quad (10)$$

In terms of the components of $|A\rangle$ and $|B\rangle$ the last expression reads

$$\langle t \rangle = \sum_{lj'l'j'} \sum_{\substack{mn \\ m'n'}} A_n^{l*} A_m^j B_{n'}^l B_{m'}^{j*} M_{nmn'm'}^{lj}, \quad (11)$$

where

$$\begin{aligned} M_{nmn'm'}^{lj} &= \sqrt{(2l+1)(2j+1)} \\ &\times \int dg \text{tr} U^{(1)}(g) \mathfrak{D}_{m'm}^{(j)}(g) \mathfrak{D}_{n'n}^{(l)*}(g) \\ &= \sqrt{(2l+1)(2j+1)} \\ &\times \sum_M \langle 1Mjm|ln\rangle \langle 1Mjm'|ln'\rangle, \end{aligned} \quad (12)$$

and the last terms in brackets are the usual Clebsch-Gordan coefficients.

The optimal strategy is the one that maximizes $\langle t \rangle$. It is tempting to introduce Lagrange multipliers λ and μ^j for the normalization constraints (5) and (8) respectively and follow the standard maximization procedure. Analytical results along this line seem hard to obtain [12]. We will, thus, try to develop a more physical picture of Eqs. 10–12 which will lead us to a stunning simplification of the problem.

Notice that Eqs. 10–12 can also be written in a compact form as

$$\langle t \rangle = \sum_{lj} \frac{\sqrt{(2l+1)(2j+1)}}{3} \langle B^j \tilde{B}^l | P_1 | A^j \tilde{A}^l \rangle, \quad (13)$$

where $|A^j \tilde{A}^l\rangle = |A^j\rangle \otimes |\tilde{A}^l\rangle$, the state $|\tilde{A}^j\rangle$ is the time reversed of $|A^j\rangle$, i.e., $\tilde{A}_m^j = (-1)^m A_{-m}^{j*}$ (and similarly for $|B^j \tilde{B}^l\rangle$ and $|\tilde{B}^l\rangle$) and P_1 is the projector over the Hilbert space of the representation of total spin $J = 1$. Our aim is to compute

$$\langle t \rangle_{\max} = \max_{AB} \langle t \rangle, \quad (14)$$

where the maximization is over all A_m^j and B_m^j subject to the normalization conditions in (5) and (8). The Schwarz inequality implies

$$\langle B^j \tilde{B}^l | P_1 | A^j \tilde{A}^l \rangle \leq \|P_1 |A^j \tilde{A}^l\rangle\| \|P_1 |B^j \tilde{B}^l\rangle\|, \quad (15)$$

where the equality holds iff

$$P_1 |A^j \tilde{A}^l\rangle = \mu^{jl} P_1 |B^j \tilde{B}^l\rangle \quad \forall j, l. \quad (16)$$

Hence, to compute $\langle t \rangle_{\max}$, we can restrict ourselves to a smaller parameter space, where $|A^j\rangle$ and $|B^j\rangle$ are constrained through (16). This is equivalent to consider the states $|A\rangle$ such that

$$A_m^j = C^j B_m^j, \quad \text{with} \quad \sum_j |C^j|^2 = 1, \quad (17)$$

i.e., we only need to consider the set of parameters $\{C^j, B_m^j\}$. This we can prove, e.g., by induction on j using (16) with $l = j + 1$ and starting with the trivial case $j = 0$ [13]. Eq. 17 is easy to understand from the physical point of view. It just tells us that, for an optimal communication, the messenger states $|A(g)\rangle$ must be as similar as possible to the states $|B(g)\rangle$ on which the measuring device projects [7]. We next substitute back in (13) to obtain

$$\langle t \rangle_{\max} = \max_{BC} \sum_{jj'} C^j M_B^{jj'} C^{j'}, \quad (18)$$

where

$$M_B^{jj'} = \frac{\sqrt{(2j+1)(2j'+1)}}{3} \langle B^j \tilde{B}^{j'} | P_1 | B^j \tilde{B}^{j'} \rangle \quad (19)$$

and the maximization is over all B_m^j and C^j subject to the normalizations (8) and (17).

Let us now discuss some properties of the matrix M_B defined by (19). We first note that M_B is tridiagonal, i.e., $M_B^{jj'} = 0$ if $|j - j'| > 1$, and symmetric. It is manifestly non-negative, i.e., $M_B^{jj'} \geq 0$ for all j, j' and, most important, it is rotationally invariant: any reference state of the form $|B'\rangle = U(g)|B\rangle$ is equally as good as $|B\rangle$.

We next compute bounds for the diagonal (M_B^{jj}) and off diagonal (M_B^{jj+1}) entries of M_B . We have

$$0 \leq M_B^{jj} = \frac{2j+1}{3} |\langle B^j \tilde{B}^j | 10 \rangle|^2 \leq \frac{2j+1}{3} \times \left| \sum_{m'} |B_{m'}^j|^2 \max_m \langle j m j - m | 10 \rangle \right|^2 = \frac{j}{j+1}, \quad (20)$$

where we have used rotational invariance to orient the (real) vector $P_1 |B^j \tilde{B}^j\rangle$ along the z ($m = 0$) axes. As for the off diagonal entries, the Schwarz inequality leads to

$$\begin{aligned} 0 \leq M_B^{jj+1} &\leq \frac{\sqrt{(2j+1)(2j+3)}}{3} \sum_{m'} |B_{m'}^j|^2 \\ &\times \sum_{m''} |\tilde{B}_{m''}^{j+1}|^2 \max_m \left(\sum_M \langle j M - m j + 1 m | 1 M \rangle^2 \right) \\ &= \sqrt{\frac{2j+1}{2j+3}}, \end{aligned} \quad (21)$$

where, actually, the sum over M in the second line is independent of m . It is straightforward to verify that the particular choice

$$|B_{\text{op}}\rangle = \sum_j \sqrt{2j+1} |j, j\rangle \quad \Leftrightarrow \quad B_{\text{op}m}^j = \delta_m^j \quad (22)$$

saturates the two upper bounds (20) and (21) *simultaneously*. Hence

$$|M_B^{jj'}| \leq M_{B_{\text{op}}}^{jj'} \equiv M_{\text{op}}^{jj'}, \quad (23)$$

for all j, j' and $|B\rangle$. The matrix $M_{\text{op}}^{jj'}$ is

$$M_{\text{op}} = \begin{pmatrix} \frac{J}{J+1} & \sqrt{\frac{2J-1}{2J+1}} & & 0 \\ \sqrt{\frac{2J-1}{2J+1}} & \ddots & \ddots & \\ & \ddots & \frac{2}{3} & \sqrt{\frac{3}{5}} \\ 0 & & \sqrt{\frac{3}{5}} & \frac{1}{2} & \sqrt{\frac{1}{3}} \\ & & & \sqrt{\frac{1}{3}} & 0 \end{pmatrix}, \quad (24)$$

where $J = N/2$ is the maximum spin of the system.

We now go back to (18) and compute $\langle t \rangle_{\text{max}}$. We first note that, $\langle t \rangle_{\text{max}} = \max_B \lambda(B)$, where $\lambda(B)$ is the

maximal eigenvalue of the matrix M_B . Since it is non-negative, Eq. 23 implies [14]

$$\langle t \rangle_{\text{max}} = \max_B \lambda(B) = \lambda(B_{\text{op}}) \equiv \lambda_{\text{op}}. \quad (25)$$

We thus have simplified the problem to that of computing λ_{op} , the maximal eigenvalues of M_{op} in (24). This can be done proceeding along the same lines as in [7, 9]. We would like to emphasize that the calculation relies on the fact that the maximal value of each entry of M_B is reached simultaneously, e.g. for the single state $|B_{\text{op}}\rangle$. This is, a priori, a rather unexpected property which, however, provides a remarkable simplification of the calculation.

The result obtained and the form of the optimal state, $|B_{\text{op}}\rangle$, agree with our physical intuition as we now briefly discuss. If Alice's state has a well defined total spin (i.e. it is an eigenstate of $\vec{J}^2$), M_{op} becomes diagonal and $\langle t \rangle_{\text{max}} = J/(J+1) = N/(N+2)$. In terms of the average error, $\langle h \rangle = 4(N+3)/(N+2)$, thus, at most ($N \rightarrow \infty$) $\langle h \rangle = 4$. In average, Bob cannot determine more than just one axes of Alice's trihedron. The structure of the state $|B_{\text{op}}\rangle$ is such that, within each irreducible representation, the determination of a single axes is optimal [2] (this is *the best* Alice could do if she only was allowed to use a single irreducible representation). At the same time, $|B_{\text{op}}\rangle$ is as different of an eigenstate of J_z as it can possibly be (if $J_z |B_{\text{op}}\rangle \propto |B_{\text{op}}\rangle$, Alice would be able to communicate *only* a single axes).

For small N , one can easily obtain analytic expressions for $\langle t \rangle_{\text{max}}$ (see table). For large N it suffices to give simple lower and upper bounds for $\langle t \rangle_{\text{max}}$. A useful upper bound is provided by the condition $\langle t \rangle_{\text{max}} \leq \max_j \sum_{j'} M_{\text{op}}^{jj'}$. A lower bound is obtained computing $\Delta = \sum_{jj'} C^j M_{\text{op}}^{jj'} C^{j'}$ for any normalized vector with components C^j . A judicious choice is $C^j \propto \sqrt{2j-1}(N/2-j)j^p$. The maximum of Δ occurs at $p \approx \sqrt[3]{3N/4}$. We obtain

$$3 - \frac{4}{N} + O(N^{-4/3}) \lesssim \langle t \rangle_{\text{max}} \lesssim 3 - \frac{4}{N} + O(N^{-2}). \quad (26)$$

It is now clear that perfect determination of the trihedron, $\langle t \rangle_{\text{max}} = 3$, is reached in the asymptotic limit, and $\langle t \rangle$ approaches three at most linearly in $1/N$. Finally, we have performed a linear fit obtaining

$$\langle t \rangle_{\text{max}} \sim 3 - \frac{4}{N} - \frac{9.4}{N^{4/3}} + \dots, \quad (27)$$

which is completely consistent with (26).

We now turn our attention to the construction of POVM's with a finite number of outcomes, as they are the only ones that can be physically realized. The main idea is stated in [10]. There, we introduced the concept of set of directions isotropically distributed. In the context of the present letter the term directions has to be generalized to elements of the group. We say that a finite set

N	2	3	5	10	50	100
$\langle t \rangle_{\max}$	$\frac{3+\sqrt{57}}{12}$	$\frac{14+\sqrt{466}}{30}$	1.6708	2.6202	2.9362	2.9707

TABLE I: Maximal value of $\langle t \rangle$ vs. the number of spins

$\{g_r\}$, $r = 1, \dots, N(J)$, of elements of $SU(2)$ is isotropically distributed up to spin J , if there exist positive weights $\{c_r\}$ such that the following orthogonality relation holds for any $j, j' \leq J$:

$$\sum_{r=1}^{N(J)} c_r \mathfrak{D}_{mn}^{(j)}(g_r) \mathfrak{D}_{m'n'}^{(j')*}(g_r) = \frac{C_J}{2j+1} \delta_m^{m'} \delta_n^{n'} \delta_j^{j'}, \quad (28)$$

where $C_J = \sum_{r=1}^{N(J)} c_r$. This discrete version of (9) is only valid up to a certain value J , the larger J is, the larger $N(J)$ must be chosen. Working along the same lines as in [10] one can show that the angular dependence on α and γ can be trivially satisfied choosing $N+1$ equidistant angles for each variable. The only non-trivial conditions concern the set $\{\beta_r\}$, which is required to satisfy $\sum_r c_r P_L(\cos \beta_r) = 0$ ($1 \leq L \leq 2J$), where P_L is the Legendre polynomial of degree L . The procedure to solve this equation is described in [10] (see also [12]). This recipe yields a finite optimal POVM for any value of N . In general, however, one can find equally optimal POVM's with a smaller number of outcomes. Ideally one would be interested in finding the minimal ones, however, as far as we are aware, the solution is not known for arbitrary J and general groups [4].

Nevertheless, the minimal POVM for the first non-trivial case of two spins is not difficult to find. Consider the simplest normalized reference state that leads to an optimal POVM

$$|B\rangle = \frac{\sqrt{3}}{2}|1, 1\rangle + \frac{1}{2}|0, 0\rangle, \quad (29)$$

It is easy to verify that the four projectors $O_r = U(g_r)|B\rangle\langle B|U^\dagger(g_r)$, with

$$\begin{aligned} \alpha_r &= (r-1)\frac{2\pi}{3}, \quad \gamma_r = \pi - \alpha_r, \quad \cos \theta_r = \frac{1}{3}, \quad r \leq 3 \\ \alpha_4 &= 0, \quad \gamma_4 = 0, \quad \cos \theta_4 = 1, \end{aligned} \quad (30)$$

satisfy the POVM condition $\sum_{r=1}^4 O_r = \mathbb{I}$. Since the Hilbert space has dimension four, the minimal number of outcomes for any measurement is also four. This measurement is therefore finite, minimal, and optimal. In fact, it is a von Neumann measurement as $O_r O_s = \delta_{rs} O_r$. Notice that the set of points (30) do not satisfy the orthogonality conditions (28) for all the m, m' values, but

it does for the relevant ones. It is the particular structure of the state (29) what enables us to construct a POVM with only four outcomes.

We conclude that it is feasible to use quantum systems to encode the orientation of a reference frame. The optimal strategy involves the use of encoding states which are remarkably simple and have a clear physical interpretation. The average error of the transmission is seen to approach zero linearly in $1/N$. Finally, we give a recipe for constructing finite optimal POVMs and present an example of a minimal one for the simple case $N = 2$.

Financial support from CICYT contract AEN99-0766 and CIRIT contracts 1998SGR-00051, 1999SGR-00097 is acknowledged.

-
- [1] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory*, North Holland, Amsterdam, 1982.
 - [2] S. Massar and S. Popescu, Phys. Rev. Lett. **74**, 1259 (1995).
 - [3] R. Derka, V. Buzek and A.K. Ekert, Phys. Rev. Lett. **80**, 1571 (1998).
 - [4] J.I. Latorre, P. Pascual and R. Tarrach, Phys. Rev. Lett. **81**, 1351 (1998).
 - [5] N. Gisin and S. Popescu, Phys. Rev. Lett. **83**, 432 (1999).
 - [6] S. Massar, Phys. Rev. **A62** 040101(R) (2000).
 - [7] E. Bagan *et al.*, Phys. Rev. Lett. **85**, 5230 (2000).
 - [8] A. Peres and P. Scudo, Phys. Rev. Lett. **86**, 4160 (2001).
 - [9] E. Bagan *et al.*, Phys. Rev. **A63**, 052309 (2001).
 - [10] E. Bagan, M. Baig and R. Muñoz-Tapia, quant-ph/0012045 [Phys. Rev. A] (in press).
 - [11] A. Acin, J.I. Latorre and P. Pascual, Phys. Rev. **A61**, 2213 (2000); S. Massar and S. Popescu, Phys. Rev. **A61**, 62303 (2000); A. Acin, R. Tarrach and G. Vidal, Phys. Rev. **A61**, 62307 (2000); A. Acin, E. Jane and G. Vidal, quant-ph/0012015.
 - [12] After our analysis was completed, we learnt about the numerical study performed by: A. Peres and P. Scudo, quant-ph/0103149. We acknowledge the correspondence with the authors.
 - [13] One can try to escape from (17) by further constraining $|B^j\rangle$ in such a way that the orthogonal complement of $P_{j+1}(\mathcal{H}^1 \otimes |B^j\rangle)$, where $\mathcal{H}^1$ is the Hilbert space of the $j = 1$ representation of $SU(2)$ and P_{j+1} is the projector over $\mathcal{H}^{j+1}$, is different from zero. Obviously, this leads to non optimal states $|A^j\rangle$ and $|B^j\rangle$.
 - [14] M. Marcus and H. Minc, *A Survey of Matrix Theory and Matrix Inequalities*, Dover Publications, Inc., New York, 1992. Statement 5.7.5, pag. 126. A simple proof is as follows. Notice that all the components of the eigenvector corresponding to the maximal eigenvalue of a non-negative matrix must have the same sign. Now, if M' and M are nonnegative matrices with $M'_{ij} \geq M_{ij} \geq 0$, then $\sum_{ij} v_i M'_{ij} v_j \geq \sum_{ij} v_i M_{ij} v_j$ for any vector with $v_i v_j \geq 0 \quad \forall i, j$.